arXiv:1608.01381v1 [math.GT] 3 Aug 2016

THE A-POLYNOMIAL 2-TUPLE OF TWISTED WHITEHEAD LINKS

ANH T. TRAN

Abstract. We compute the A-polynomial 2-tuple of twisted Whitehead links. As applications, we determine canonical components of twisted Whitehead links and give a formula for the volume of twisted Whitehead link cone-manifolds.

1. Introduction

The A-polynomial of a knot in the 3-sphere S^3 was introduced by Cooper, Culler, Gillet, Long and Shalen [CCGLS] in the 90's. It describes the $SL_2(\mathbb{C})$-character variety of the knot complement as viewed from the boundary torus. The A-polynomial carries a lot of information about the topology of the knot. For example, it distinguishes the unknot from other knots [BoZ, DG] and the sides of the Newton polygon of the A-polynomial give rise to incompressible surfaces in the knot complement [CCGLS].

The A-polynomial was generalized to links by Zhang [Zh] about ten years later. For an m-component link in S^3, Zhang defined a polynomial m-tuple link invariant called the A-polynomial m-tuple. The A-polynomial 1-tuple of a knot is nothing but the A-polynomial defined in [CCGLS]. The A-polynomial m-tuple also caries important information about the topology of the link. For example, it can be used to construct concrete examples of hyperbolic link manifolds with non-integral traces [Zh].

Finding an explicit formula for the A-polynomial is a challenging problem. So far, the A-polynomial has been computed for a few classes of knots including two-bridge knots $C(2n, p)$ (with $1 \leq p \leq 5$) in Conway's notation [HS, Pe, Ma, HL], $(-2, 3, 2n+1)$-pretzel knots [TY, GM]. It should be noted that $C(2n, p)$ is the double twist knot $J(2n, -p)$ in the notation of [HS]. Moreover, $C(2n, 1)$ is the torus knot $T(2, 2n+1)$ and $C(2n, 2)$ is known as a twist knot. A cabling formula for the A-polynomial of a cable knot in S^3 has recently been given in [NZ]. Using this formula, Ni and Zhang [NZ] has computed the A-polynomial of an iterated torus knot explicitly.

In this paper we will compute the A-polynomial 2-tuple for a family of 2-component links called twisted Whitehead links. As applications, we will determine canonical components of twisted Whitehead links and give a formula for the volume of twisted Whitehead link cone-manifolds. For $k \geq 0$, the k-twisted Whitehead link W_k is the 2-component link depicted in Figure 1. Note that W_0 is the torus link $T(2, 4)$ and W_1 is the Whitehead link. Moreover, W_k is the two-bridge link $C(2, k, 2)$ in Conway's notation and is $\mathfrak{b}(4k+4, 2k+1)$ in Schubert's notation. These links are all hyperbolic except for W_0.

The A-polynomial 2-tuple of the twisted Whitehead link W_k is a polynomial 2-tuple $[A_1(M, L), A_2(M, L)]$ given as follows.

2000 *Mathematics Subject Classification.* Primary 57M27, Secondary 57M25.

Key words and phrases. canonical component, cone-manifold, hyperbolic volume, the A-polynomial, twisted Whitehead link, two-bridge link.



FIGURE 1. The k-twisted Whitehead link W_k.

Theorem 1.1. *If $k = 2n-1$ then $A_1(M,L) = (L-1)F(M,L)$ where*

$$\begin{aligned} F(M,L) &= \sum_{i=0}^{n} \left\{ \binom{n+1+i}{2i+1} - \binom{n-1+i}{2i+1} \right\} (M-M^{-1})^{2i} \left(\frac{L-1}{L+1}\right)^{2i} \\ &\quad + \sum_{i=0}^{n-1} \binom{n+i}{2i+1} (M+M^{-1})(M-M^{-1})^{2i+1} \left(\frac{L-1}{L+1}\right)^{2i+1}. \end{aligned}$$

If $k = 2n$ then $A_1(M,L) = (LM^2-1)G(M,L)$ where

$$\begin{aligned} G(M,L) &= \sum_{i=0}^{n} \left\{ \binom{n+1+i}{2i+1} + \binom{n+i}{2i+1} \right\} (M-M^{-1})^{2i+1} \left(\frac{LM^2-1}{LM^2+1}\right)^{2i+1} \\ &\quad + \sum_{i=0}^{n} \binom{n+i}{2i} (M+M^{-1})(M-M^{-1})^{2i} \left(\frac{LM^2-1}{LM^2+1}\right)^{2i}. \end{aligned}$$

In both cases we have $A_1(M,L) = A_2(M,L)$.

It should be noted that in the definition of the A-polynomial m-tuple of an m-component link, we discard the irreducible component of the $SL_2(\mathbb{C})$-character variety containing only characters of reducible representations and therefore consider only the nonabelian $SL_2(\mathbb{C})$-character variety of the link group.

For a link $\mathcal{L}$ in S^3, let $E_{\mathcal{L}} = S^3 \setminus \mathcal{L}$ denote the link exterior. The link group of $\mathcal{L}$ is defined to be the fundamental group $\pi_1(E_{\mathcal{L}})$ of $E_{\mathcal{L}}$. It is known that the link group of the twisted Whitehead link W_k has a standard two-generator presentation of a two-bridge link group $\pi_1(E_{W_k}) = \langle a, b \mid aw = wa \rangle$, where a, b are meridians depicted in Figure 1 and w is a word in the letters a, b. More precisely, we have $w = (bab^{-1}a^{-1})^n a (a^{-1}b^{-1}ab)^n$ if $k = 2n-1$ and $w = (bab^{-1}a^{-1})^n bab (a^{-1}b^{-1}ab)^n$ if $k = 2n$.

For a presentation $\rho : \pi_1(E_{W_k}) \to SL_2(\mathbb{C})$ we let x, y, z denote the traces of the images of a, b, ab respectively. We also let v denote the trace of the image of $bab^{-1}a^{-1}$. Explicitly, we have $v = x^2 + y^2 + z^2 - xyz - 2$. It was shown in [Tr1] that the nonabelian $SL_2(\mathbb{C})$-character variety of W_k has exactly $\lfloor \frac{k}{2} \rfloor$ irreducible components. For a hyperbolic link, there are distinguished components of the $SL_2(\mathbb{C})$-character variety called the canonical components. They contain important information about the hyperbolic structure of the link. By using the formula of the A-polynomial in Theorem 1.1, we can determine the canonical component of the hyperbolic twisted Whitehead link W_k (with $k \geq 1$) as follows.

Theorem 1.2. *If $k = 2n-1$ then the canonical component of W_k is the zero set of the polynomial $(xy - vz)S_{n-1}(v) - (xy - 2z)S_{n-2}(v)$.*

If $k = 2n$ then the canonical component of W_k is the zero set of the polynomial $zS_n(v) - (xy - z)S_{n-1}(v)$.

Here the $S_k(v)$ are the Chebychev polynomials of the second kind defined by $S_0(v) = 1$, $S_1(v) = v$ and $S_k(v) = vS_{k-1}(v) - S_{k-2}(v)$ for all integers k. Explicitly, we have

$$S_k(v) = \sum_{0 \le i \le k/2} (-1)^i \binom{k-i}{k-2i} v^{k-2i}$$

for $k \ge 0$, $S_k(v) = -S_{-k-2}(v)$ for $k \le -2$, and $S_{-1}(v) = 0$.

For a hyperbolic link $\mathcal{L} \subset S^3$, let $\rho_{\rm hol}$ be a holonomy representation of $\pi_1(E_{\mathcal{L}})$ into $PSL_2(\mathbb{C})$. Thurston [Th] showed that $\rho_{\rm hol}$ can be deformed into an m-parameter family $\{\rho_{\alpha_1,\cdots,\alpha_m}\}$ of representations to give a corresponding family $\{E_{\mathcal{L}}(\alpha_1, \cdots, \alpha_m)\}$ of singular complete hyperbolic manifolds, where m is the number of components of $\mathcal{L}$. In this paper we will consider only the case where all of α_i's are equal to a single parameter α. In which case we also denote $E_{\mathcal{L}}(\alpha_1, \cdots, \alpha_m)$ by $E_{\mathcal{L}}(\alpha)$. These α's and $E_{\mathcal{L}}(\alpha)$'s are called the cone-angles and hyperbolic cone-manifolds of $\mathcal{L}$, respectively.

We consider the complete hyperbolic structure on a link complement as the cone-manifold structure of cone-angle zero. It is known that for a two-bridge knot or link $\mathcal{L}$ there exists an angle $\alpha_{\mathcal{L}} \in [\frac{2\pi}{3}, \pi)$ such that $E_{\mathcal{L}}(\alpha)$ is hyperbolic for $\alpha \in (0, \alpha_{\mathcal{L}})$, Euclidean for $\alpha = \alpha_{\mathcal{L}}$, and spherical for $\alpha \in (\alpha_{\mathcal{L}}, \pi)$ [HLM, Ko1, Po, PW]. Explicit volume formulas for hyperbolic cone-manifolds of knots were known for 4_1 [HLM, Ko1, Ko2, MR], 5_2 [Me1], twist knots [HMP] and two-bridge knots $C(2n, 3)$ [HL]. Recently, the volume of double twist knot cone-manifolds has been computed in [Tr3]. We should remark that a formula for the volume of the cone-manifold of $C(2n, 4)$ has just been given in [HLMR1]. However, with an appropriate change of variables, this formula has already obtained in [Tr3], since $C(2n, 4)$ is the double twist knot $J(2n, -4)$.

In this paper we are interested in hyperbolic cone-manifolds of links. Explicit volume formulas for hyperbolic cone-manifolds of links have been only known for 5_1^2 [MV], 6_2^2 [Me2], 6_3^2 [DMM] and 7_3^2 [HLMR2]. From the proof of Theorem 1.1, we can compute the volume of the hyperbolic cone-manifold of the twisted Whitehead link W_k as follows.

Let

$$R_{W_k}(s, z) = \begin{cases} x^2 S_{n-1}(v) - zS_n(v) - (x^2 - z)S_{n-2}(v) & \text{if } k = 2n-1 \\ zS_n(v) - (x^2 - z)S_{n-1}(v) & \text{if } k = 2n, \end{cases}$$

where $x = s + s^{-1}$ and $v = 2x^2 + z^2 - x^2 z - 2$.

Theorem 1.3. *For $\alpha \in (0, \alpha_{W_k})$ we have*

$$\operatorname{Vol} E_{W_k}(\alpha) = \int_{\alpha}^{\pi} 2 \log \left| \frac{z - (s^{-2} + 1)}{z - (s^2 + 1)} \right| d\omega$$

where $s = e^{i\omega/2}$ and z (with $\operatorname{Im} z \ge 0$) satisfy $R_{W_k}(s, z) = 0$.

Note that the above volume formula for the hyperbolic cone-manifold $E_{W_k}(\alpha)$ depends on the choice of a root z of $R_{W_k}(e^{i\omega/2}, z) = 0$. In practice, we choose the root z which gives the maximal volume.

As a direct consequence of Theorem 1.3, we obtain the following.

Corollary 1.4. *The hyperbolic volume of the r-fold cyclic covering over the twisted Whitehead link W_k, with $r \geq 3$, is given by the following formula*

$$r\mathrm{Vol}\, E_{W_k}(\frac{2\pi}{r}) = r\int_{\frac{2\pi}{r}}^{\pi} 2\log\left|\frac{z-(s^{-2}+1)}{z-(s^2+1)}\right| d\omega$$

where $s = e^{i\omega/2}$ and z (with $Im\, z \geq 0$) satisfy $R_{W_k}(s,z) = 0$.

Remark 1.5. The link 7_3^2 is the twisted Whitehead link W_3. With an appropriate change of variables, we obtain the volume formula for the hyperbolic cone-manifold of 7_3^2 in [HLMR2, Theorem 1.1] by taking $k = 3$ in Theorem 1.3.

The paper is organized as follows. In Section 2 we review the definition of the A-polynomial m-tuple of an m-component link in S^3. In Section 3 we review the nonabelian $SL_2(\mathbb{C})$-representations of a two-bridge link and compute them explicitly for twisted Whitehead links. We also give proofs of Theorems 1.1–1.3 in Section 3.

2. The A-polynomial

2.0.1. *Character varieties.* The set of characters of representations of a finitely generated group G into $SL_2(\mathbb{C})$ is known to be a complex algebraic set, called the character variety of G and denoted by $\chi(G)$ (see [CS, LM]). For a manifold Y we also use $\chi(Y)$ to denote $\chi(\pi_1(Y))$. Suppose $G = \mathbb{Z}^2$, the free abelian group with 2 generators. Every pair of generators μ, λ will define an isomorphism between $\chi(G)$ and $(\mathbb{C}^*)^2/\tau$, where $(\mathbb{C}^*)^2$ is the set of non-zero complex pairs (M, L) and $\tau : (\mathbb{C}^*)^2 \to (\mathbb{C}^*)^2$ is the involution defined by $\tau(M, L) := (M^{-1}, L^{-1})$, as follows. Every representation is conjugate to an upper diagonal one, with M and L being the upper left entry of μ and λ respectively. The isomorphism does not change if we replace (μ, λ) with (μ^{-1}, λ^{-1}).

2.0.2. *The A-polynomial.* Suppose $\mathcal{L} = K_1 \sqcup \cdots \sqcup K_m$ be an m-component link in S^3. Let $E_{\mathcal{L}} = S^3 \setminus \mathcal{L}$ be the link exterior and $T_1, \ldots, T_m$ the boundary tori of $E_{\mathcal{L}}$ corresponding to $K_1, \ldots, K_m$ respectively. Each T_i is a torus whose fundamental group is free abelian of rank two. An orientation of K_i will define a unique pair of an oriented meridian μ_i and an oriented longitude λ_i such that the linking number between the longitude λ_i and the knot K_i is 0. The pair provides an identification of $\chi(\pi_1(T_i))$ and $(\mathbb{C}^*)_i^2/\tau_i$, where $(\mathbb{C}^*)_i^2$ is the set of non-zero complex pairs (M_i, L_i) and τ_i is the involution $\tau(M_i, L_i) = (M_i^{-1}, L_i^{-1})$, which actually does not depend on the orientation of K_i.

The inclusion $T_i \hookrightarrow E_{\mathcal{L}}$ induces the restriction map

$$\rho_i : \chi(E_{\mathcal{L}}) \longrightarrow \chi(T_i) \equiv (\mathbb{C}^*)_i^2/\tau_i.$$

For each $\gamma \in \pi_1(E_{\mathcal{L}})$ let f_γ be the regular function on the character variety $\chi(E_{\mathcal{L}})$ defined by $f_\gamma(\chi_\rho) = (\chi_\rho(\gamma))^2 - 4$, where χ_ρ denotes the character of a representation $\rho : \pi_1(E_{\mathcal{L}}) \to SL_2(\mathbb{C})$. Let $\chi_i(X)$ be the subvariety of $\chi(X)$ defined by $f_{\mu_j} = 0,\ f_{\lambda_j} = 0$ for all $j \neq i$. Let Z_i be the image of $\chi_i(X)$ under ρ_i and $\hat{Z}_i \subset (\mathbb{C}^*)_i^2$ the lift of Z_i under the projection $(\mathbb{C}^*)_i^2 \to (\mathbb{C}^*)_i^2/\tau_i$. It is known that the Zariski closure of $\hat{Z}_i \subset (\mathbb{C}^*)_i^2 \subset \mathbb{C}_i^2$ in $\mathbb{C}_i^2$ is an algebraic set consisting of components of dimension 0 or 1 [Zh]. The union of all the 1-dimension components is defined by a single polynomial $A_i' \in \mathbb{Z}[M_i, L_i]$ whose coefficients are co-prime. Note that A_i' is defined up to ± 1. It is also known that A_i' always contains the factor $L_i - 1$ coming from the characters of reducible representations, hence $A_i' = (L_i - 1)A_i$

for some $A_i \in \mathbb{Z}[M_i, L_i]$. As in [Zh], we will call $[A_1(M_1, L_1), \cdots, A_m(M_m, L_m)]$ the A-polynomial m-tuple of $\mathcal{L}$. For brevity, we also write $A_i(M, L)$ for $A_i(M_i, L_i)$. We refer the reader to [Zh] for properties of the A-polynomial m-tuple.

Recall that the Newton polygon of a two-variable polynomial $\sum a_{ij} M^i L^j$ is the convex hull in $\mathbb{R}^2$ of the set $\{(i, j) : a_{ij} \neq 0\}$. The slope of a side of the Newton polygon is called a boundary slope of the polygon. The following proposition is useful for determining canonical components of hyperbolic links.

Proposition 2.1. [Zh, Theorem 3(2)] *Suppose $\mathcal{L} \subset S^3$ is a hyperbolic m-component link. For each $j = 1, \cdots, m$, the factor $A_j^{\text{can}}(M, L)$ of the A-polynomial $A_j(M, L)$ corresponding to a canonical component has an irreducible factor whose Newton polygon has at least two distinct boundary slopes. In particular, this irreducible factor contains at least 3 monomials in M, L.*

3. Proofs of Theorems 1.1–1.3

In this section we review the nonabelian $SL_2(\mathbb{C})$-representations of a two-bridge link from [Ri] and compute them explicitly for twisted Whitehead links. Finally, we give proofs of Theorems 1.1–1.3.

3.1. **Two-bridge links.** Two-bridge links are those links admitting a projection with only two maxima and two minima. The double branched cover of S^3 along a two-bridge link is a lens space $L(2p, q)$, which is obtained by doing a $2p/q$ surgery on the unknot. Such a two-bridge link is denoted by $\mathfrak{b}(2p, q)$. Here q is an odd integer co-prime with $2p$ and $2p > |q| \geq 1$. It is known that $\mathfrak{b}(2p', q')$ is ambient isotopic to $\mathfrak{b}(2p, q)$ if and only if $p' = p$ and $q' \equiv q^{\pm 1} \pmod{2p}$, see e.g. [BuZ]. The link group of the two-bridge link $\mathfrak{b}(2p, q)$ has a standard two-generator presentation $\pi_1(E_{\mathfrak{b}(2p,q)}) = \langle a, b \mid wa = aw \rangle$ where a, b are meridians, $w = b^{\varepsilon_1} a^{\varepsilon_2} \cdots a^{\varepsilon_{2p-2}} b^{\varepsilon_{2p-1}}$ and $\varepsilon_i = (-1)^{\lfloor iq/(2p) \rfloor}$ for $1 \leq i \leq 2p-1$. Note that $\varepsilon_i = \varepsilon_{2p-i}$.

We now study representations of link groups into $SL_2(\mathbb{C})$. A representation is called nonabelian if its image is a nonabelian subgroup of $SL_2(\mathbb{C})$. Let $\mathcal{L} = \mathfrak{b}(2p, q)$. Suppose $\rho : E_{\mathcal{L}} \to SL_2(\mathbb{C})$ is a nonabelian representation. Up to conjugation, we may assume that

$$\rho(a) = \begin{bmatrix} s_1 & 1 \\ 0 & s_1^{-1} \end{bmatrix} \quad \text{and} \quad \rho(b) = \begin{bmatrix} s_2 & 0 \\ u & s_2^{-1} \end{bmatrix}$$

where $(u, s_1, s_2) \in \mathbb{C}^3$ satisfies the matrix equation $\rho(aw) = \rho(wa)$.

For any word r, we write $\rho(r) = \begin{bmatrix} r_{11} & r_{12} \\ r_{21} & r_{22} \end{bmatrix}$. By induction on the word length, we can show that r_{21} is a multiple of u in $\mathbb{C}[u, s_1^{\pm 1}, s_2^{\pm 1}]$. Hence we can write $r_{21} = u r'_{21}$ for some $r'_{21} \in \mathbb{C}[u, s_1^{\pm 1}, s_2^{\pm 1}]$. A word is said to be *palindromic* if it reads the same backward or forward. By Lemma 1 in [Ri] we have

$$r_{22} - r_{11} + (s_1 - s_1^{-1}) r_{12} = (s_2 - s_2^{-1}) r'_{21}$$

for any palindromic word r of odd length.

The matrix equation $\rho(aw) = \rho(wa)$ is easily seen to be equivalent to the two equations $u w'_{21} = 0$ and $w_{22} - w_{11} + (s_1 - s_1^{-1}) w_{12} = (s_2 - s_2^{-1}) w'_{21}$. Since w is a palindromic word of odd length, we conclude that the matrix equation $\rho(aw) = \rho(wa)$ is equivalent to a single equation $w'_{21} = 0$. The polynomial w'_{21} is called the Riley polynomial of a two-bridge link, and it determines the nonabelian representations of the link.

Remark 3.1. Other approaches, using character varieties and skein modules, to the Riley polynomial of a two-bridge link can be found in [Qa, Tr2, LT].

3.2. **Chebyshev polynomials.** Recall that the $S_k(v)$ are the Chebychev polynomials of the second kind defined by $S_0(v) = 1$, $S_1(v) = v$ and $S_k(v) = vS_{k-1}(v) - S_{k-2}(v)$ for all integers k. The following results are elementary, see e.g. [Tr3].

Lemma 3.2. *For any integer k we have*

$$S_k^2(v) + S_{k-1}^2(v) - vS_k(v)S_{k-1}(v) = 1.$$

Lemma 3.3. *Suppose $V \in SL_2(\mathbb{C})$ and $v = \operatorname{tr} V$. For any integer k we have*

$$V^k = S_{k-1}(v)V - S_{k-2}(v)\mathbf{1}$$

where $\mathbf{1}$ denotes the 2×2 identity matrix.

We will need the following lemma in the proof of Theorem 1.1.

Lemma 3.4. *Suppose $v = 2 + q$. For $k \geq 0$ we have*

$$S_k(v) = \sum_{i=0}^{k} \binom{k+1+i}{2i+1} q^i. \tag{3.1}$$

Proof. We use induction on $k \geq 0$. The cases $k = 0, 1$ are clear. Suppose $k \geq 2$ and (3.1) holds true for $k-2$ and $k-1$. Since $S_k(v) = vS_{k-1}(v) - S_{k-2}(v)$, we have

$$\begin{aligned} S_k(v) &= (2+q)\sum_{i=0}^{k-1} \binom{k+i}{2i+1} q^i - \sum_{i=0}^{k-2} \binom{k-1+i}{2i+1} q^i \\ &= \sum_{i=0}^{k} \left\{ 2\binom{k+i}{2i+1} + \binom{k+i-1}{2i-1} - \binom{k-1+i}{2i+1} \right\} q^i. \end{aligned}$$

It remains to show the following identity

$$\binom{k+1+i}{2i+1} = 2\binom{k+i}{2i+1} + \binom{k+i-1}{2i-1} - \binom{k-1+i}{2i+1}.$$

This follows by applying the equality $\binom{c}{d} + \binom{c}{d+1} = \binom{c+1}{d+1}$ three times. □

3.3. **Twisted Whitehead links.** In this subsection we compute nonabelian representations of twisted Whitehead links explicitly.

We first consider the case of W_{2n-1}. Recall that the link group of W_{2n-1} has a presentation $\pi_1(E_{W_{2n-1}}) = \langle a, b \mid aw = wa \rangle$, where a, b are meridians depicted in Figure 1 and $w = (bab^{-1}a^{-1})^n a (a^{-1}b^{-1}ab)^n$.

Suppose $\rho : \pi_1(E_{W_{2n-1}}) \to SL_2(\mathbb{C})$ is a nonabelian representation. Up to conjugation, we may assume that

$$\rho(a) = \begin{bmatrix} s_1 & 1 \\ 0 & s_1^{-1} \end{bmatrix} \quad \text{and} \quad \rho(b) = \begin{bmatrix} s_2 & 0 \\ u & s_2^{-1} \end{bmatrix}$$

where $(u, s_1, s_2) \in \mathbb{C}^3$ satisfies the matrix equation $\rho(aw) = \rho(wa)$.

Recall from the Introduction that $x = \operatorname{tr}\rho(a) = s_1 + s_1^{-1}$, $y = \operatorname{tr}\rho(b) = s_2 + s_2^{-1}$, $z = \operatorname{tr}\rho(ab) = u + s_1 s_2 + s_1^{-1}s_2^{-1}$ and

$$v = \operatorname{tr}\rho(bab^{-1}a^{-1}) = u(u + s_1 s_2 + s_1^{-1}s_2^{-1} - s_1 s_2^{-1} - s_1^{-1}s_2) + 2.$$

Let $c = bab^{-1}a^{-1}$ and $d = a^{-1}b^{-1}ab$. We have

$$\rho(c) = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix} \qquad \text{and} \qquad \rho(d) = \begin{bmatrix} d_{11} & d_{12} \\ d_{21} & d_{22} \end{bmatrix}$$

where

$$\begin{aligned}
c_{11} &= 1 - s_1^{-1}s_2u, \\
c_{12} &= -s_1 + s_1s_2^2 + s_2u, \\
c_{21} &= u(-s_1^{-2}s_2^{-1} + s_2^{-1} - s_1^{-1}u), \\
c_{22} &= 1 + (s_1^{-1}s_2^{-1} - s_1s_2^{-1} + s_1s_2)u + u^2, \\
d_{11} &= 1 + (s_1^{-1}s_2^{-1} - s_1^{-1}s_2 + s_1s_2)u + u^2, \\
d_{12} &= s_1^{-1}s_2^{-2} - s_1^{-1} + s_2^{-1}u, \\
d_{21} &= u(s_2 - s_1^2s_2 - s_1u), \\
d_{22} &= 1 - s_1s_2^{-1}u.
\end{aligned}$$

Since $v = \operatorname{tr}\rho(c) = \operatorname{tr}\rho(d)$, by Lemma 3.3 we have

$$\begin{aligned}
\rho(c^n) &= \begin{bmatrix} c_{11}S_{n-1}(v) - S_{n-2}(v) & c_{12}S_{n-1}(v) \\ c_{21}S_{n-1}(v) & c_{22}S_{n-1}(v) - S_{n-2}(v) \end{bmatrix}, \\
\rho(d^n) &= \begin{bmatrix} d_{11}S_{n-1}(v) - S_{n-2}(v) & d_{12}S_{n-1}(v) \\ d_{21}S_{n-1}(v) & d_{22}S_{n-1}(v) - S_{n-2}(v) \end{bmatrix}.
\end{aligned}$$

This implies that $\rho(w) = \begin{bmatrix} w_{11} & * \\ w_{21} & * \end{bmatrix} = \rho(c^nad^n)$ with

$$\begin{aligned}
w_{11} &= (c_{12}d_{21}s_1^{-1} + c_{11}d_{21} + c_{11}d_{11}s_1)S_{n-1}^2(v) + s_1S_{n-2}^2(v) \\
&\quad - (d_{21} + c_{11}s_1 + d_{11}s_1)S_{n-1}(v)S_{n-2}(v), \\
w_{21} &= \big[(c_{22}d_{21}s_1^{-1} + c_{21}d_{21} + c_{21}d_{11}s_1)S_{n-1}(v) - (d_{21}s_1^{-1} + c_{21}s_1)S_{n-2}(v)\big]S_{n-1}(v).
\end{aligned}$$

By direct calculations we have

$$\begin{aligned}
c_{22}d_{21}s_1^{-1} + c_{21}d_{21} + c_{21}d_{11}s_1 &= u(xy - vz), \\
d_{21}s_1^{-1} + c_{21}s_1 &= u(xy - 2z).
\end{aligned}$$

Hence $w_{21} = uw'_{21}$ where

$$w'_{21}(W_{2n-1}) = \big((xy - vz)S_{n-1}(v) - (xy - 2z)S_{n-2}(v)\big)S_{n-1}(v),$$

which is the Riley polynomial of W_{2n-1}.

Similarly, the Riley polynomial of W_{2n} is given by the following formula

$$w'_{21}(W_{2n}) = \big(zS_n(v) - (xy - z)S_{n-1}(v)\big)\big(S_n(v) - S_{n-1}(v)\big).$$

Remark 3.5. The above formulas for the Riley polynomials of twisted Whitehead links were already obtained in [Tr1] using character varieties. It was also shown in [Tr1] that the Riley polynomial $w'_{21}(W_{2n-1}) \in \mathbb{C}[x, y, z]$ is factored into exactly n irreducible factors $(xy-vz)S_{n-1}(v)-(xy-2z)S_{n-2}(v)$ and $v-2\cos\frac{j\pi}{n}$ $(1 \le j \le n-1)$. Similarly, $w'_{21}(W_{2n}) \in \mathbb{C}[x, y, z]$ is factored into exactly $n + 1$ irreducible factors $zS_n(v) - (xy - z)S_{n-1}(v)$ and

$v-2\cos\frac{(2j-1)\pi}{2n+1}$ $(1\le j\le n)$. Note that

$$S_{n-1}(v)=\prod_{j=1}^{n-1}\big(v-2\cos\frac{j\pi}{n}\big)\quad\text{and}\quad S_n(v)-S_{n-1}(v)=\prod_{j=1}^{n}\big(v-2\cos\frac{(2j-1)\pi}{2n+1}\big).$$

3.4. **Proof of Theorem 1.1.** We first consider the case of W_{2n-1}. The canonical longitudes corresponding to the meridians a and b are respectively $\lambda_a=wa^{-1}$ and $\lambda_b=\overline{w}b^{-1}$, where $\overline{w}$ is the word obtained from w by exchanging a and b. Precisely, we have $\overline{w}=(aba^{-1}b^{-1})^n b(b^{-1}a^{-1}ba)^n$.

Suppose $\rho:\pi_1(E_{W_{2n-1}})\to SL_2(\mathbb{C})$ is a nonabelian representation. With the same notations as in the previous subsection, we have

$$\rho(w)=\begin{bmatrix} w_{11} & * \\ 0 & (w_{11})^{-1}\end{bmatrix}\quad\text{and}\quad \rho(\overline{w})=\begin{bmatrix} \overline{w}_{11} & 0 \\ * & (\overline{w}_{11})^{-1}\end{bmatrix}.$$

Proposition 3.6. *On $v-2\cos\frac{j\pi}{n}=0$ we have*

$$w_{11}=s_1\quad\textit{and}\quad \overline{w}_{11}=s_2.$$

Proof. Recall from the previous subsection that

$$\begin{aligned} w_{11} &= (c_{12}d_{21}s_1^{-1}+c_{11}d_{21}+c_{11}d_{11}s_1)S_{n-1}^2(v)+s_1S_{n-2}^2(v)\\ &\quad-(d_{21}+c_{11}s_1+d_{11}s_1)S_{n-1}(v)S_{n-2}(v).\end{aligned}$$

If $v=2\cos\frac{j\pi}{n}$ then $S_{n-1}(v)=0$ then $S_{n-2}(v)=\pm1$. Hence $w_{11}=s_1$.

The proof for $\overline{w}_{11}=s_2$ is similar. □

Proposition 3.7. *On $(xy-vz)S_{n-1}(v)-(xy-2z)S_{n-2}(v)=0$ we have*

$$w_{11}=\frac{-1-s_2^2+s_1s_2z}{s_1+s_1s_2^2-s_2z}\quad\textit{and}\quad \overline{w}_{11}=\frac{-1-s_1^2+s_1s_2z}{s_2+s_1^2s_2-s_1z}.$$

Proof. By Lemma 3.2 we have $S_k^2(v)+S_{k-1}^2(v)-vS_k(v)S_{k-1}(v)=1$ for all integers k. Hence, on $(xy-vz)S_{n-1}(v)-(xy-2z)S_{n-2}(v)=0$ we have

$$S_{n-1}^2(v)=\big(1-rv+r^2\big)^{-1}$$

where $r=(xy-vz)/(xy-2z)$. By a direct calculation we have

$$\begin{aligned} w_{11} &= \big[(c_{12}d_{21}s_1^{-1}+c_{11}d_{21}+c_{11}d_{11}s_1)+s_1r^2-(d_{21}+c_{11}s_1+d_{11}s_1)r\big]\big(1-rv+r^2\big)^{-1}\\ &= \frac{s_1s_2(s_1^2s_2+s_1u-s_2)}{s_1^2-s_1s_2u-1}=\frac{-1-s_2^2+s_1s_2z}{s_1+s_1s_2^2-s_2z}.\end{aligned}$$

The proof for $\overline{w}_{11}$ is similar. □

We now prove Theorem 1.1 for W_{2n-1}. Let $\rho(\lambda_a)=\begin{bmatrix} L_a & * \\ 0 & L_a^{-1}\end{bmatrix}$. Since $\lambda_a=wa^{-1}$, we have $L_a=w_{11}s_1^{-1}$. On

$$C_j:=\Big(\{v-2\cos\frac{j\pi}{n}=0\}\cap\{s_2^2=(\overline{w}_{11})^2=1\}\Big)$$

by Proposition 3.6 we have $w_{11}=s_1$. Hence $L_a-1=0$ on C_j for all $1\le j\le n-1$.

By Proposition 3.7, on

$$C:=\Big(\{(xy-vz)S_{n-1}(v)-(xy-2z)S_{n-2}(v)=0\}\cap\{s_2^2=(\overline{w}_{11})^2=1\}\Big)$$

we have $w_{11} = \frac{-1-s_2^2+s_1s_2z}{s_1+s_1s_2^2-s_2z}$. Note that on C, we have $s_2^2 = 1$ implies $(\overline{w}_{11})^2 = 1$.

With $s_2 = 1$ we then have $w_{11} = \frac{-2+s_1z}{2s_1-z}$. Hence $z = \frac{2(1+s_1w_{11})}{s_1+w_{11}}$. Note that $v = 2+(z-x)^2$. Let $z - x = t$. Then

$$t = \frac{2(1+s_1w_{11})}{s_1+w_{11}} - (s_1+s_1^{-1}) = (s_1-s_1^{-1})\big(\frac{w_{11}-s_1}{w_{11}+s_1}\big) = (s_1-s_1^{-1})\Big(\frac{L_a-1}{L_a+1}\Big)$$

and $v = 2+t^2$. Since $y = s_2+s_2^{-1} = 2$ and $vS_{n-1}(v) = S_n(v)+S_{n-2}(v)$ we have

$$\begin{aligned} 0 &= 2xS_{n-1}(v) - zS_n(v) - (2x-z)S_{n-2}(v) \\ &= 2xS_{n-1}(v) - (x+t)S_n(v) - (x-t)S_{n-2}(v) \\ &= -xt^2S_{n-1}(v) - t\big(S_n(v) - S_{n-2}(v)\big). \end{aligned}$$

If $t = 0$ then $w_{11} = \frac{-2+s_1x}{2s_1-x} = s_1$. In this case we have $L_a = 1$. We now consider the case $xtS_{n-1}(v) + S_n(v) - S_{n-2}(v) = 0$. Then, by Lemma 3.4 we have

$$\begin{aligned} 0 &= \sum_{i=0}^{n}\left\{\binom{n+1+i}{2i+1} - \binom{n-1+i}{2i+1}\right\}t^{2i} + x\sum_{i=0}^{n-1}\binom{n+i}{2i+1}t^{2i+1} \\ &= \sum_{i=0}^{n}\left\{\binom{n+1+i}{2i+1} - \binom{n-1+i}{2i+1}\right\}(s_1-s_1^{-1})^{2i}\left(\frac{L_a-1}{L_a+1}\right)^{2i} \\ &\quad + \sum_{i=0}^{n-1}\binom{n+i}{2i+1}(s_1+s_1^{-1})(s_1-s_1^{-1})^{2i+1}\left(\frac{L_a-1}{L_a+1}\right)^{2i+1}. \end{aligned}$$

The last expression in the above equalities is exactly the polynomial $F(M_a, L_a)$, with $M_a = s_1$, defined in Theorem 1.1. Hence, with $s_2 = 1$, we have $(L_a-1)F(M_a, L_a) = 0$ on C. Similarly, with $s_2 = -1$, we obtain the same equation $(L_a-1)F(M_a, L_a) = 0$ on C. This proves the formula of the A-polynomial for the component of W_{2n-1} corresponding to the meridian a. The one for the component corresponding b is exactly the same.

For W_{2n} and its component corresponding to a, we have $L_aM_a^2 - 1 = 0$ on

$$D_j := \Big(\{v - 2\cos\frac{(2j-1)\pi}{2n+1} = 0\} \cap \{s_2^2 = (\overline{w}_{11})^2 = 1\}\Big)$$

for $1 \le j \le n$. Moreover, on

$$D := \Big(\{zS_n(v) - (xy-z)S_{n-1}(v) = 0\} \cap \{s_2^2 = (\overline{w}_{11})^2 = 1\}\Big)$$

we have $G(M_a, L_a) = 0$ where G is the polynomial defined in Theorem 1.1. The same formulas can be obtained for the component of W_{2n} corresponding to b.

This completes the proof of Theorem 1.1.

3.5. **Proof of Theorem 1.2.** We make use of Proposition 2.1 which implies that for each $j = 1, 2$ the factor $A_j^{\text{can}}(M, L)$ of the A-polynomial $A_j(L, M)$ corresponding to a canonical component of W_k (with $k \ge 1$) has an irreducible factor containing at least 3 monomials in M, L. This irreducible factor cannot be $L-1$ or LM^2-1. Hence, from the proof of Theorem 1.1 we conclude that the canonical components of W_{2n-1} and W_{2n} are respectively the zero sets of the polynomials $(xy-vz)S_{n-1}(v) - (xy-2z)S_{n-2}(v)$ and $zS_n(v) - (xy-z)S_{n-1}(v)$. This completes the proof of Theorem 1.2.

3.6. **Proof of Theorem 1.3.** Recall that $E_{W_k}(\alpha)$ is the cone-manifold of W_k with cone angles $\alpha_1 = \alpha_2 = \alpha$. There exists an angle $\alpha_{W_k} \in [\frac{2\pi}{3}, \pi)$ such that $E_{W_k}(\alpha)$ is hyperbolic for $\alpha \in (0, \alpha_{W_k})$, Euclidean for $\alpha = \alpha_{W_k}$, and spherical for $\alpha \in (\alpha_{W_k}, \pi)$.

We first consider the case of W_{2n-1}. By Theorem 1.2 the canonical component of W_{2n-1} is determined by $(xy - vz)S_{n-1}(v) - (xy - 2z)S_{n-2}(v) = 0$. Moreover, by Proposition 3.7, on this component we have

$$w_{11} = \frac{-1 - s_2^2 + s_1 s_2 z}{s_1 + s_1 s_2^2 - s_2 z}.$$

Here we use the same notations as in the previous subsections.

For $\alpha \in (0, \alpha_{W_{2n-1}})$, by the Schlafli formula we have

$$\operatorname{Vol} E_{W_{2n-1}}(\alpha) = \int_\alpha^\pi 2 \log |w_{11}| \, d\omega = \int_\alpha^\pi 2 \log \left| \frac{-1 - s_2^2 + s_1 s_2 z}{s_1 + s_1 s_2^2 - s_2 z} \right| d\omega$$

where $s_1 = s_2 = s = e^{i\omega/2}$ and z (with $\left|\frac{-1-s_2^2+s_1s_2z}{s_1+s_1s_2^2-s_2z}\right| \geq 1$) satisfy

$$(xy - vz)S_{n-1}(v) - (xy - 2z)S_{n-2}(v) = 0.$$

We refer the reader to [DMM] for the volume formula of hyperbolic cone-manifolds of links using the Schlafli formula.

Note that $\left|\frac{-1-s_2^2+s_1s_2z}{s_1+s_1s_2^2-s_2z}\right| = \left|\frac{z-(s^{-2}+1)}{z-(s^2+1)}\right| \geq 1$ is equivalent to $\operatorname{Im} z \geq 0$. Since $x = y = s + s^{-1}$ and $v = x^2 + y^2 + z^2 - xyz - 2 = 2x^2 + z^2 - x^2 z - 2$, the proof of Theorem 1.3 for W_{2n-1} is complete. The proof for W_{2n} is similar.

Acknowledgements

The author has been partially supported by a grant from the Simons Foundation (#354595 to Anh Tran).

References

[BoZ] S. Boyer and X. Zhang, *Every non-trivial knot in S^3 has non-trivial A-polynomial*, Proc. Amer. Math. Soc. **133** (2005), no. 9, 2813–2815 (electronic).

[BuZ] G. Burde and H. Zieschang, *Knots*, de Gruyter Stud. Math., vol. 5, de Gruyter, Berlin, 2003.

[CCGLS] D. Cooper, M. Culler, H. Gillet, D. Long and P. Shalen, *Plane curves associated to character varieties of 3-manifolds*, Invent. Math. **118** (1994), pp. 47–84.

[CS] M. Culler and P. Shalen, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math. (2) **117** (1983), no. 1, 109–146.

[DG] N. Dunfield and S. Garoufalidis, *Non-triviality of the A-polynomial for knots in S^3*, Algebr. Geom. Topol. **4** (2004), 1145–1153 (electronic).

[DMM] D. Derevnin, A. Mednykh and M. Mulazzani, *Volumes for twist link cone-manifolds*, Bol. Soc. Mat. Mexicana (3) **10** (2004), Special Issue, 129–145.

[GM] S. Garoufalidis and T. Mattman, *The A-polynomial of the $(-2, 3, 3+2n)$-pretzel knots*, New York J. Math. **17** (2011), 269–279.

[HL] J. Ham and J. Lee, *An explicit formula for the A-polynomial of the knot with Conway's notation $C(2n, 3)$*, preprint 2016, arXiv:1601.05860.

[HLM] H. Hilden, M. Lozano, and J. Montesinos-Amilibia, *Volumes and Chern-Simons invariants of cyclic coverings over rational knots*, in Topology and Teichmüller spaces (Katinkulta, 1995), pages 31–55. World Sci. Publ., River Edge, NJ, 1996.

[HLMR1] J. Ham, J. Lee, A. Mednykh and A. Rasskazov, *On the volume and the Chern-Simons invariant for the* 2*-bridge knot orbifolds*, preprint 2016, arXiv:1607.08044.

[HLMR2] J. Ham, J. Lee, A. Mednykh and A. Rasskazov, *An explicit volume formula for the link* $7_3^2(\alpha,\alpha)$ *cone-manifolds*, preprint 2016, arXiv:1607.08047.

[HMP] J.-Y. Ham, A. Mednykh, and V. Petrov, *Identities and volumes of the hyperbolic twist knot cone-manifolds*, J. Knot Theory Ramifications **2**3 (2014) 1450064.

[HS] J. Hoste and P. Shanahan, *A formula for the A-polynomial of twist knots*, J. Knot Theory Ramifications **14** (2005), 91–100.

[Ko1] S. Kojima, *Deformations of hyperbolic 3-cone-manifolds*, J. Differential Geom. **49** (1998) 469–516.

[Ko2] S. Kojima, *Hyperbolic 3-manifolds singular along knots*, Chaos Solitons Fractals **9** (1998) 765–777.

[LM] A. Lubotzky and A. Magid, *Varieties of representations of finitely generated groups*, Memoirs of the AMS **336** (1985).

[LT] T. Le and A. Tran, *The Kauffman bracket skein module of two-bridge links*, Proc. Amer. Math. Soc. **142** (2014), no. 3, 1045–1056.

[Ma] Daniel Mathews, *An explicit formula for the A-polynomial of twist knots*, J. Knot Theory Ramifications **23** (2014), no. 9, 1450044, 5 pp.

[Me1] A. Mednykh, *The volumes of cone-manifolds and polyhedra* http://mathlab.snu.ac.kr/$\tilde{\text{t}}$op/workshop01.pdf, 2007. Lecture Notes, Seoul National University.

[Me2] A. Mednykh, *Trigonometric identities and geometrical inequalities for links and knots*, Proceedings of the Third Asian Mathematical Conference, 2000 (Diliman), 352–368, World Sci. Publ., River Edge, NJ, 2002.

[MR] A. Mednykh and A. Rasskazov, *Volumes and degeneration of cone-structures on the figure-eight knot*, Tokyo J. Math. **29** (2006) 445–464.

[MV] A Mednykh and A. Vesnin, *On the volume of hyperbolic Whitehead link cone-manifolds*, Geometry and analysis. Sci. Ser. A Math. Sci. (N.S.) **8** (2002), 1–11.

[NZ] Y. Ni and X. Zhang, *Detection of knots and a cabling formula for A-polynomials*, preprint 2014, arXiv:1411.0353.

[Pe] K. Petersen, *A-polynomials of a family of two-bridge knots*, New York J. Math. **21** (2015), 847–881.

[Po] J. Porti, *Spherical cone structures on 2-bridge knots and links*, Kobe J. Math. **21** (2004) 61–70.

[PW] J. Porti and H. Weiss, *Deforming Euclidean cone 3-manifolds*, Geom. Topol. **11** (2007) 1507–1538.

[Qa] K. Qazaqzeh, *The character variety of a family of one-relator groups*, Internat. J. Math. **23** (2012), no. 1, 1250015, 12 pp.

[Ri] R. Riley, *Algebra for Heckoid groups*, Trans. Amer. Math. Soc. **334** (1992), 389–409.

[Th] W. Thurston, *The geometry and topology of 3-manifolds*, http://library.msri.org/books/gt3m, 1977/78. Lecture Notes, Princeton University.

[Tr1] A. Tran, *Character varieties of* $(-2,2m+1,2n)$*-pretzel links and twisted Whitehead links*, J. Knot Theory Ramifications **25** (2016), no. 2, 1650007, 16 pp.

[Tr2] A. Tran, *The universal character ring of the* $(-2,2m+1,2n)$*-pretzel link*, Internat. J. Math. **24** (2013), no. 8, 1350063, 13 pp.

[Tr3] A. Tran, *Volumes of double twist knot cone-manifolds*, preprint 2015, arXiv:1512.08165.

[TY] N. Tamura and Y. Yokota, *A formula for the A-polynomials of* $(-2, 3, 1+2n)$*-pretzel knots*, Tokyo J. Math. **27** (2004), no. 1, 263–273.

[Zh] X. Zhang, *The A-polynomial n-tuple of a link and hyperbolic 3-manifolds with non-integral traces*, J. Knot Theory Ramifications **15** (2006), 279–287.

Department of Mathematical Sciences, The University of Texas at Dallas, Richardson, TX 75080, USA

E-mail address: att140830@utdallas.edu